

MILLING OATS VARIETY TRIAL


ABSTRACT:

Since 2016, Gateway Research Organization (GRO) has collaborated with the Prairie Oat Growers Association (POGA) to conduct an in-depth evaluation of 11 approved oat milling varieties, focusing on their performance and beta-glucan content. There are two locations in north central Alberta - with Gateway Research Organization (GRO) at (Westlock, and with Smoky Applied Research & Demonstration Association (SARDA) at Falher. This ongoing initiative aims to provide valuable insights into how different oat varieties and regional growing conditions influence both yield and the functional attributes associated with beta-glucan levels. Over the years, the study has consistently highlighted notable varietal differences in yield and beta-glucan content in the two regions. Nine years of research were successfully completed through to 2024, and POGA has agreed to continue its support for an additional three years, covering the 2025, 2026, and 2027 growing seasons. This ongoing commitment will ensure that the research continues to advance and support the oat industry.

PROJECT BACKGROUND:

Oats are a beneficial rotational crop for producers to use, as they contribute to soil fertility and pest management. They provide disease and insect breaks for wheat, barley, and canola. Their rapid establishment and growth provide excellent weed suppression. Oats also work well as a “catch crop” for taking up and storing excess nitrogen, and the straw provides a nutrient source for the following year’s crop. The straw also protects against soil erosion and contributes to an increase in the soil’s organic matter content (Campbell et al., 1991). Well-planned management and appropriate selection of varieties make oats a profitable crop due to their low input requirements and favourable effects on succeeding crops in a rotation. Test weight is the most commonly used measure of grain quality. Growers who plan to sell oat grain typically aim for varieties with high test weights. However, in the processing industry, other properties, such as the solubility and viscosity of beta-glucan, are often more important. Research has shown that oat beta β -glucan can help lower blood cholesterol, regulate blood sugar, and improve insulin response, which in turn may reduce the risk of heart disease and support diabetes prevention (Wang and Ellis, 2014).

Oat acreage in Alberta declined in the early 2010s but has been on a steady upward path since 2018. After a slight dip in 2023, the trend recovered in 2024 and continues to grow into 2025. Statistics Canada reports that the total area planted with oats increased by roughly 3.3 percent, reaching 1.213 million hectares compared to 2024.



Source: Government of Canada, Canadian Grain Commission (2025, November 4). Quality of Western Canadian Oats in 2025. <https://www.grainscanada.gc.ca/en/grain-research/grain-harvest-export-quality/oats/2025/harvest-quality-report-oats-2025.html>

ALBERTA MILLING OAT TRIAL CONTINUED

This graph shows how the area seeded with different oat varieties in western Canada has changed from 2020 to 2025. CS Camden started off as the most popular choice, covering about 33 percent of the area in 2020, but its popularity steadily dropped to around 17 percent by 2025. Meanwhile, CDC Arborg gained a lot of ground, starting at just 3 percent and growing to nearly 17 percent, catching up to CS Camden by 2025. **AC Morgan stayed relatively steady, fluctuating around 18 to 20 percent throughout the years.** Summit saw a slow decline from about 16 percent down to just above 10 percent. CDC Endure started out almost unnoticed but gradually grew to cover about 8 percent of the area by 2025. CDC Anson appeared in 2024 and quickly jumped to nearly 10 percent in 2025. Overall, the graph tells a story of changing preferences, with some long-time favourites giving way to newer varieties that are becoming more popular.

To use the “Heart Healthy” label, a product must contain at least 4% beta β -glucan. The β -glucan content in oats may vary with changes in growing conditions (Perez Herrera et al., 2016). **In Alberta, the most commonly grown oat variety is AC Morgan, but it often struggles to reach the necessary beta-glucan levels, except in drier years.** Additionally, AC Morgan does not have crown rust resistance. Oat farmers in the region need a variety that can reliably match or exceed AC Morgan’s yields while also providing higher beta-glucan content and a strong disease resistance package. Since 2015, Grain Millers Canada Corporation has supported these variety trials, aiming to find oat types that help Alberta producers access the milling market more consistently.

Helping producers discover oat varieties that perform better in their region is an important step toward keeping their farms profitable. This trial will give farmers in Alberta useful insights, allowing them to grow oats that not only yield more but also offer added health benefits, like higher levels of beta-glucan.

OBJECTIVE:

The objective of the milling oat trial is to investigate how genotype and growing conditions affect the yield and β -glucan content of milling oat varieties in Northern Alberta.

METHODOLOGY:

This study was conducted at two sites, **Westlock and Falher**. At each location, a uniform field was selected for the trial, and soil testing was conducted to guide nutrient management. Fertilizer was applied according to the soil test recommendations to ensure optimal growing conditions. Seeding rates for each variety were calculated based on the 1000-kernel weight, target plant density (300 plants/m²), and germination percentage. Planting was carried out using a small 6-row Fabro seeder. Each plot measured 1.37 meters in width and 7 meters in length, with four replications included for statistical reliability. Herbicides were applied as needed to maintain weed-free plots. Harvesting was carried out using a Zurn 150 at Westlock and a Wintersteiger plot harvester at Falher. Total yields from each plot were recorded, and clean sub-samples were taken to create 500-gram composite samples, which were submitted for beta-glucan analysis.

2025 SOIL INFORMATION

	Nitrogen (lbs/ac)	Phosphorous (lbs/ac)	Potassium (lbs/ac)	Sulphur (lbs/ac)	pH (0-14)	CEC (meq/100g)	Organic Matter (%)
Westlock	20	36	216	25	6.3	20.1	6.8
Peace Region	20	56	256	30	4.8	19.2	4.7

Before the start of the season, POGA reviewed the entry list and made a few updates. **They decided to remove CDC Ruffian and AAC Neville from the study and added CDC Hank and CDC Fetch in their place.**

Table: Agronomic Information

	Westlock	Falher
Legal Land Description	SW17-61-26-W4	SW29-77-20-W5
Seeded:	02-May-25	15-May-25
Seeding Depth:	1 ¼ inch	¾ inch
Soil Moisture:	Excellent	Good
Soil Temp:	12°C	13.1°C
Rainfall:	194.3 mm	102.5 mm
Fertilizer:	105.9N-35P-60K-20S	123N-40P-20K-30S
Pre-emergence:	None	Conquer II + Roundup Weathermax @ 200 ml/ac + 1000 ml/ac on May 22, 2025
In-crop Application:	Buctril M @ 400 ml/ac on June 3, 2025	Stellar XL @ 405 ml/ac on June 18, 2025
Desiccant:	Reglone Ion @ 1 L/ac on Sept. 4, 2025	None
Harvested:	September 16, 2025	September 21, 2025

RESULTS AND DISCUSSION:

In 2025 at Westlock, the site averaged 143.3 bu/ac, and variety selection had a significant impact on yield according to the ANOVA test, which showed clear statistical differences among entries. **CDC Anson** and **AAC Anthony** were the top performers, both yielding just over 152 bu/ac and finishing well above the site average and the check, **AC Morgan**. Other varieties such as **CS Camden**, **CDC Endure**, and **AAC Fetch** also performed slightly above the check.

At the Falher site, the average yield was 157.6 bu/ac, with strong performance across varieties. **AAC Anthony** led the trial at 167.7 bu/ac, followed closely by **CDC Byer** and **AAC Wesley**, all delivering excellent yields. However, the ANOVA test indicated that differences among varieties were not statistically significant at this location, meaning yields were more uniform despite the numerical spread.

ALBERTA MILLING OAT TRIAL CONTINUED

2025 YIELD COMPARISON

		Westlock			Peace Region		
Trt No.	Treatment Name	% of AC Morgan	Yield (bu/ac)		% of AC Morgan	Yield (bu/ac)	
1	AC Morgan	100	143.0	b	100	158.4	a
2	CS Camden	102	145.5	b	100	157.9	a
3	CDC Arborg	100	143.0	b	95	150.4	a
4	CDC Endure	101	144.3	b	96	152.6	a
5	AAC Douglas	98	140.8	b	98	154.7	a
6	CDC Byer	97	138.5	b	104	164.5	a
7	CDC Anson	107	152.5	a	96	151.4	a
8	CDC Hank	98	140.3	b	101	159.7	a
9	AAC Wesley	92	132.0	c	101	160.6	a
10	AAC Anthony	106	152.0	a	106	167.7	a
11	AAC Fetch	101	143.8	b	98	155.4	a

Means followed by same letter or symbol do not significantly differ (P=.05, Student-Newman-Keuls).
Mean comparisons performed only when AOV Treatment P(F) is significant at mean comparison OSL.

Test weight is a vital parameter in assessing the milling quality of grain, particularly oats. A higher test weight typically correlates with better processing characteristics, reduced waste and improved end product quality, making it an essential metric for both producers and processors in the oat industry.



MORE RESULTS FROM THE POGA TRIAL - GRO, WESTLOCK - 2025

Trt	Treatment	Height		Lodging		Test Weight				1000 KWT	
No.	Name	(cm)		1 (erect) - 9 (flat)		lbs/bu		kg/HL		(g)	
1	AC Morgan	107.8	b	1.5	bc	44.688	ab	55.1	ab	41.4	bcd
2	CS Camden	100.5	c	1.0	c	44.838	ab	55.3	ab	39.3	de
3	CDC Arborg	110.3	ab	2.0	b	45.088	a	55.6	a	40.1	cde
4	CDC Endure	111.8	a	1.0	c	43.298	bcd	53.4	bcd	42.3	b
5	AAC Douglas	101.3	c	1.5	bc	43.525	abc	53.7	abc	41.9	bc
6	CDC Byer	101.3	c	1.5	bc	45.0	ab	55.6	a	40.0	cde
7	CDC Anson	91.5	e	1.0	c	44.49	ab	54.9	ab	40.3	b-e
8	CDC Hank	102.0	c	1.5	bc	42.878	cd	52.9	cd	42.5	b
9	AAC Wesley	91.0	e	1.0	c	44.0	abc	54.3	abc	39.0	e
10	AAC Anthony	109.3	ab	1.8	bc	42.128	d	52.0	d	46.2	a
11	AAC Fetch	96.3	d	3.5	a	44.153	abc	54.5	abc	38.9	e
LSD P=.05 (% mean diff)		2.68 (3%)		0.60 (38%)		1.02 (3%)		1.25 (3%)		1.55 (4%)	
Standard Deviation		1.85		0.41		0.70		0.87		1.08	
CV		1.82		26.39		1.6		1.6		2.62	

Means followed by same letter or symbol do not significantly differ (P=.05, Student-Newman-Keuls).

Mean comparisons performed only when AOV Treatment P(F) is significant at mean comparison OSL.

MORE RESULTS FROM THE POGA TRIAL - SARDA, PEACE REGION - 2025

Trt	Treatment	Height		Lodging		Test Weight				1000 KWT	
No.	Name	(cm)		1 (erect) - 9 (flat)		lbs/bu		kg/HL		(g)	
1	AC Morgan	75.3	ab	1	na	43.2	ab	53.3	ab	35.5	bc
2	CS Camden	71.8	ab	1	na	41.1	c	50.7	c	34.1	bc
3	CDC Arborg	77.4	a	1	na	43.2	ab	53.2	ab	34.5	bc
4	CDC Endure	77.9	a	1	na	42.8	abc	52.8	abc	36.5	abc
5	AAC Douglas	73.4	ab	1	na	41.4	bc	51.1	bc	34.6	bc
6	CDC Byer	70.2	ab	1	na	44.0	a	54.3	a	34.9	bc
7	CDC Anson	62.4	c	1	na	42.2	bc	52.0	bc	34.7	bc
8	CDC Hank	75.9	ab	1	na	43.0	ab	53.1	ab	37.3	ab
9	AAC Wesley	66.8	bc	1	na	41.2	bc	50.9	bc	33.7	c
10	AAC Anthony	72.3	ab	1	na	41.7	bc	51.4	bc	38.6	a
11	AAC Fetch	71.8	ab	1	na	42.0	bc	51.8	bc	34.0	bc
LSD P=.05 (% mean diff)		5.81 (8%)		.		1.22 (3%)		1.50 (3%)		1.97 (6%)	
Standard Deviation		4.01		0		0.84		1.04		1.36	
CV		5.55		0		1.98		1.98		3.86	

Means followed by same letter or symbol do not significantly differ (P=.05, Student-Newman-Keuls).


Mean comparisons performed only when AOV Treatment P(F) is significant at mean comparison OSL.

ALBERTA MILLING OAT TRIAL CONTINUED

The Falher site received less precipitation (about 40% of their regional average) during the growing season, and that likely contributed to the shorter crop height and lower test weights observed there. When moisture is limited, the grain filling period is often shorter or less effective, which can result in lighter kernels and reduced bushel weight. That helps explain why the Falher averaged 42.3 lbs per bushel, compared to 44.0 lbs per bushel at Westlock. At Westlock, we also saw lodging in AAC Fetch, whereas under the drier conditions in the Peace Region no lodging was observed in any variety. CDC Byer stood out at both sites, recording test weights above the site averages and showing strong consistency across environments. Overall, the results highlight how closely plant growth, standability, and grain quality are tied to seasonal moisture and overall growing conditions.

BETA-GLUCAN TEST RESULTS FROM POGA TRIAL 2025

Trt No.	Treatment Name	Westlock (GRO)		Falher (SARDA)	
		Hull Percentage (%)	Flour BG (% db)	Hull Percentage (%)	Flour BG (% db)
1	AC Morgan	24.5	5.2	31.3	4.2
2	CS Camden	26.7	5.5	31.0	4.8
3	CDC Arborg	26.1	5.5	29.3	4.9
4	CDC Endure	21.3	5.2	26.7	5.3
5	AAC Douglas	23.3	5.1	32.4	5.0
6	CDC Byer	25.2	5.6	27.0	4.9
7	CDC Anson	23.5	5.9	27.2	5.8
8	CDC Hank	24.0	5.5	26.9	5.3
9	AAC Wesley	21.9	5.3	28.0	4.7
10	AAC Anthony	25.9	5.1	33.8	4.4
11	AAC Fetch	28.1	4.9	32.3	4.7



All the oat varieties in the trial exceeded the industry's minimum standard of 4% beta-glucan, confirming they meet the expected nutritional benchmark. **CDC Anson** was the standout variety, consistently showing the highest beta-glucan content in both Westlock at 5.9% and the Peace Region at 5.8%. CDC Hank also performed reliably at both locations. Interestingly, **AC Morgan** recorded beta-glucan levels above 5% at the Westlock site, which is unusual given that data from the past nine years has consistently shown lower values for this variety. The reason for this unexpected result is not certain. One possibility is that favourable growing conditions during critical stages of grain development led to a temporary increase in beta-glucan content. Alternatively, a sample mix-up may have occurred, either during preparation at the facility or during laboratory analysis. Further investigation is recommended to determine whether this observation reflects a true increase in nutritional quality or an anomaly in sampling or measurement.

OVERALL TRIAL SUMMARY - TOP 3 VARIETIES FOR BETA-GLUCAN FROM 2016-2025

Crop Year	Westlock		
2025	CDC Anson	CDC Byer	CDC Hank
2024	CDC Anson	CDC Endure	CDC Byer
2023	AAC Douglas	CDC Anson	AAC Wesley
2022	CDC Endure	OT3112	AAC Douglas
2021	OT3112	CDC Endure	CDC Skye
2020	OT3112	CDC Endure	CDC Skye
2019	CDC Endure	CDC Arborg	AC Morgan
2018	CDC Endure	CDC Arborg	Triactor
2017	CS Camden	Akina	CDC Ruffian
2016	CDC Seabiscuit	CDC Ruffian	CDC Orin
Crop Year	Falher		
2025	CDC Anson	CDC Endure	CDC Hank
2024	CDC Anson	CS Camden	AAC Douglas
2023	CDC Anson	CDC Endure	OT 6024
2022	CDC Endure	OT 6024	CDC Arborg
2021	OT3112	CDC Endure	CDC Skye
2020	CDC Skye	OT3112	CDC Endure
2019	CDC Seabiscuit	CDC Arborg	CS Camden
2018	Triactor	AC Morgan	CDC Endure
2017	CDC Ruffian	CS Camden	CDC Orin
2016	CDC Ruffian	AC Morgan	CDC Seabiscuit

ALBERTA MILLING OAT TRIAL CONTINUED

OVERALL TRIAL SUMMARY - YIELD FROM 2016 TO 2025 AT WESTLOCK

Milling Oats	Yield % of AC Morgan	Overall Avg. Yield (bu/ac)	2025	2024	2023	2022	2021	2020	2019	2018	2017	2016
			Yield (bu/ac)									
AC Morgan	100	199	143	179	257	192	161	203	243	226	212	178
CS Camden	99	198	146	186	257	189	150	211	241	206	226	167
CDC Arborg	101	202	143	186	263	198	150	208	244	221	-	-
CDC Endure	100	199	144	190	252	195	143	194	249	226	-	-
CDC Anson - OT3112	95	189	153	180	254	195	140	213	-	-	-	-
AAC Douglas	92	183	141	173	261	193	148	-	-	-	-	-
AAC Wesley	92	183	132	171	230	199	-	-	-	-	-	-
CDC Byer	85	169	139	199	-	-	-	-	-	-	-	-
AAC Anthony	81	163	152	173	-	-	-	-	-	-	-	-
CDC Hank	70	140	140	-	-	-	-	-	-	-	-	-
AAC Fetch	72	144	144	-	-	-	-	-	-	-	-	-
CDC Ruffian	103	205	-	184	239	208	147	206	219	207	245	193
AAC Neville	85	169	-	169	-	-	-	-	-	-	-	-
Kalio	96	191	-	-	252	180	141	-	-	-	-	-
ORe Level 50	103	205	-	-	227	182	-	-	-	-	-	-
OT 6024	109	217	-	-	241	193	-	-	-	-	-	-
AC Summit	95	189	-	-	-	-	121	178	245	203	217	167
CDC Skye	94	188	-	-	-	-	115	211	237	-	-	-
ORe3541M	58	115	-	-	-	-	115	-	-	-	-	-
CDC Seabiscuit	105	210	-	-	-	-	-	205	237	212	208	189
ORe3542M	100	199	-	-	-	-	-	183	214	201	-	-
CDC Norseman	104	208	-	-	-	-	-	190	222	213	-	-
Triactor	106	212	-	-	-	-	-	-	238	229	208	172
Akina	103	206	-	-	-	-	-	-	-	221	222	176
CDC Orrin	101	202	-	-	-	-	-	-	-	218	221	168
Souris	88	175	-	-	-	-	-	-	-	-	194	155
Kara	100	199	-	-	-	-	-	-	-	-	222	175

OVERALL TRIAL SUMMARY - YIELD FROM 2016 TO 2025 AT FALHER

Milling Oats	Yield % of AC Morgan	Overall Avg. Yield (bu/ac)	2025	2024	2023	2022	2021	2020	2019	2018	2017	2016
			Yield (bu/ac)									
AC Morgan	100	189	158	184	187	235	20	211	224	252	220	203
CS Camden	98	186	158	173	192	265	29	183	232	217	226	190
CDC Arborg	98	187	150	179	194	269	28	199	236	237	-	-
CDC Endure	96	182	153	182	184	240	25	206	225	243	-	-
CDC Anson - OT3112	85	162	151	172	177	268	23	180	-	-	-	-
AAC Douglas	83	157	155	173	185	254	20	-	-	-	-	-
AAC Wesley	106	200	161	185	190	266	-	-	-	-	-	-
CDC Byer	91	173	165	181	-	-	-	-	-	-	-	-
AAC Anthony	93	176	168	185	-	-	-	-	-	-	-	-
CDC Hank	84	160	160	-	-	-	-	-	-	-	-	-
AAC Fetch	82	155	155	-	-	-	-	-	-	-	-	-
CDC Ruffian	104	197	-	188	188	259	21	207	203	241	249	218
AAC Neville	94	178	-	178	-	-	-	-	-	-	-	-
Kalio	80	151	-	-	184	248	22	-	-	-	-	-
ORe Level 50	106	200	-	-	181	219	-	-	-	-	-	-
OT 6024	118	223	-	-	196	250	-	-	-	-	-	-
AC Summit	91	173	-	-	-	-	19	181	227	228	210	173
CDC Skye	75	143	-	-	-	-	20	196	213	-	-	-
ORe3541M	14	27	-	-	-	-	27	-	-	-	-	-
CDC Seabiscuit	117	221	-	-	-	-	-	196	240	242	224	203
ORe3542M	110	209	-	-	-	-	-	197	205	225	-	-
CDC Norseman	113	214	-	-	-	-	-	190	214	238	-	-
Triactor	120	227	-	-	-	-	-	-	224	256	240	189
Akina	114	215	-	-	-	-	-	-	-	242	214	190
CDC Orrin	112	211	-	-	-	-	-	-	-	239	227	168
Souris	95	180	-	-	-	-	-	-	-	-	191	169
Kara	110	208	-	-	-	-	-	-	-	-	226	190

ALBERTA MILLING OAT TRIAL CONTINUED

OVERALL SUMMARY OF BETA-GLUCAN CONTENT FROM 2016 - 2025

Milling Oats	Average		2024		2023		2022		2021		2020		2019		2018		2017		2016	
	WL	PR	WL	PR	WL	PR	WL	PR	WL	PR	WL	PR	WL	PR	WL	PR	WL	PR	WL	PR
AC Morgan	3.80	3.85	4.75	4.86	3.21	3.36	3.3	3.7	3.5	3.5	3.9	3.8	3.9	3.7	3.9	3.4	3.8	4.2	3.9	4.1
CS Camden	4.32	4.47	5.28	5.59	4.2	4.61	3.8	4.2	4	4	4.7	4.3	4.4	5.2	4.4	3.8	4.4	4.6	3.7	3.9
CDC Ruffian	3.67	3.76	4.59	4.41	3.51	3.3	3.6	5.1	3.3	3.9	4.3	3.5	3.6	3.7	3.6	2.7	3.8	3.9	2.7	3.3
CDC Arborg	4.38	4.39	5.24	4.90	4.25	4.41	4.2	5.5	3.8	4.2	4.6	3.6	4.2	4.3	4.4	3.8	-	-	-	-
CDC Endure	4.75	4.92	5.31	5.41	4.52	5.04	4.9	6	4.1	4.5	5.2	4.6	4.5	4.7	4.7	4.2	-	-	-	-
CDC Anson	5.43	5.31	6.08	6.13	5.19	5.12	4.9	5.4	4.9	5.1	6.1	4.8	-	-	-	-	-	-	-	-
AC Douglas	4.87	4.71	5.14	5.43	5.85	4.11	4.8	5.2	3.7	4.1	-	-	-	-	-	-	-	-	-	-
AAC Wesley	4.59	4.93	4.90	5.14	4.88	4.35	4	5.3	-	-	-	-	-	-	-	-	-	-	-	-
CDC Byer	5.30	5.17	5.30	5.17	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
AAC Anthony	5.06	5.00	5.06	5.00	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
AAC Neville	4.52	4.88	4.52	4.88	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Kalio	4.01	3.67	-	-	3.84	3.11	4.6	4.1	3.6	3.8	-	-	-	-	-	-	-	-	-	-
ORE Level 50	4.43	4.28	-	-	4.16	3.35	4.7	5.2	-	-	-	-	-	-	-	-	-	-	-	-
OT 6024	4.53	5.37	-	-	4.76	4.83	4.3	5.9	-	-	-	-	-	-	-	-	-	-	-	-
AC Summit	4.12	4.05	-	-	-	-	-	-	3.4	3.4	4.8	4.5	4.3	4.6	4.3	3.7	4.3	4.4	3.6	3.7
CDC Skye	4.47	4.73	-	-	-	-	-	-	4	4.2	4.9	5	4.5	5	-	-	-	-	-	-
ORE 3541M	3.60	3.80	-	-	-	-	-	-	3.6	3.8	-	-	-	-	-	-	-	-	-	-
CDC	4.36	4.04	-	-	-	-	-	-	-	-	4.6	4	4.5	4.2	4.4	3.7	4.6	4.6	3.7	3.7
Seabiscuit	4.07	3.83	-	-	-	-	-	-	-	-	4.4	3.8	3.8	4.2	4	3.5	-	-	-	-
ORE3542M	4.67	4.27	-	-	-	-	-	-	-	-	4.8	4.6	4.7	4.4	4.5	3.8	-	-	-	-
CDC	4.10	4.13	-	-	-	-	-	-	-	-	-	-	4.1	4.3	4.4	4	4.4	4.5	3.5	3.7
Norseman	4.53	4.20	-	-	-	-	-	-	-	-	-	-	-	-	4.8	4	5	4.9	3.8	3.7
Triactor	3.90	3.70	-	-	-	-	-	-	-	-	-	-	-	-	4.1	3.4	4.4	4	3.2	3.7
Akina	4.25	4.40	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4.9	4.4	3.6	4.4
CDC Orrin	3.95	4.35	-	-	-	-	-	-	-	-	-	-	-	-	-	-	4.3	5	3.6	3.7
Souris	3.40	3.90	-	-	-	-	-	-	-	-	-	-	-	-	-	-	3.9	4.3	2.9	3.5
Kara																				
CDC Minstrel																				

In conclusion, weather conditions once again played a pivotal role in shaping trial outcomes this year. Both sites experienced rainfall below the long-term averages for their regions, with the Peace Region facing particularly severe drought, receiving less than 40 percent of its typical precipitation. By late June, the Peace Region recorded the lowest soil moisture ratings in the province, and a state of agricultural disaster was declared in the Municipal District of Smoky River at the beginning of August. This lack of moisture explains why plants were shorter, exhibited little to no lodging, and why average yields were substantially lower than in previous years.

This comprehensive study underscores the potential of modern genetics to deliver strong performance in both yield and quality. **However, it is important to note that not all newer varieties will consistently outperform older ones. Careful research and selection remain essential when choosing the right variety.**

Overall, both cultivar selection and location significantly influence crop yield and beta-glucan levels. Environmental factors continue to play a critical role in determining a variety's productivity and quality traits, highlighting the ongoing need for research to optimize performance under diverse conditions.

ACKNOWLEDGEMENTS:

We would like to thank the Prairie Oat Growers Association (POGA) and Grain Millers Canada for their full financial assistance. Special thanks to Paul Richter (Oat Breeder at General Mills) for contributions to lab analysis for this trial.

We would like to thank Alliance Seed, Canterra Seeds, Canada Seed Depot, FP Genetics, and SeCan for their generous seed donations for the trial. This information is presented with the understanding that no product discrimination is intended, and neither endorsement of any variety/product mentioned, nor criticism of the named variety/products is implied.

We would also like to express our gratitude to SARDA for their continued support and partnership on this trial for so many years.

